

Theoretical Computations of Transition Metal NMR Chemical Shifts

F.T. Mauschick, M. Bühl

published in

Modern Methods and Algorithms of Quantum Chemistry,
J. Grotendorst (Ed.), John von Neumann Institute for Computing,
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 2, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

Theoretical Computations of Transition Metal NMR Chemical Shifts

Frank Thomas Mauschick, Michael Bühl

*Max-Planck-Institut für Kohlenforschung, Kaiser-Wilhelm-Platz 1,
45470 Mülheim an der Ruhr, Germany*

Density functional theory (DFT) is now well established for the computation of NMR properties including chemical shifts of transition-metal compounds [1]. In most cases the B3LYP combination of density functionals is better suited than others to describe the shifts of the metal nuclei, $\delta(^{95}\text{Mo})$ being the only exception so far [1]. We have now extended the series of examined nuclei to include ^{49}Ti [2] and ^{99}Ru chemical shifts [3], both of which are well reproduced with B3LYP (Figure 1) .




Figure 1: Computed (GIAO-B3LYP) versus experimental transition-metal chemical shifts; left: ^{49}Ti , right: ^{99}Ru . (dotted: ideal slope = 1, dashed: linear fit)

References

- [1] M. Bühl, M. Kaupp, V. G. Malkin, O. L. Malkina, *J. Comput. Chem.* **1999**, 20, 91.
- [2] F. T. Mauschick, M. Bühl, unpublished results.
- [3] M. Bühl, S. Gaemers, C. J. Elsevier, submitted for publication.